

Mode-Locking Hysteresis in the Globally Coupled Model of Charge-Density Waves

Afshin Montakhab, J. M. Carlson, and Jeremy Levy

Department of Physics, University of California, Santa Barbara, CA 93106-9530

Abstract

We study the response of a recently proposed global coupling model of charge-density waves to a joint ac+dc drive. The model is the standard Fukuyama-Lee-Rice model with an additional global coupling term to account for interaction with (uncondensed) quasi-particles. We find that traditional mode-locking is accompanied by hysteresis on mode-locked steps. The experimentally observed asymmetry in the size of hysteresis is reproduced and explained by drawing attention to the importance of the pinning force. For large values of global coupling, increased rigidity leads to vanishing of subharmonic steps consistent with experiments.

PACS numbers: 72.15.Nj, 05.45.+b

Charge-density wave (CDW) conductors have been established as ideal systems in which to study the dynamics of many degrees of freedom in the presence of disorder [1]. CDWs display a variety of nonlinear nonequilibrium phenomena, including dynamical phase transitions [2] and mode-locking [3]. In the presence of an applied dc-field E_{dc} , above a threshold E_{th} , the CDW depins and carries an average current J_{cdw} which varies continuously with E_{dc} . A similar behavior is observed at the mode-locking transitions, where in the presence of joint ac+dc drive the “narrow-band noise” frequency, $\omega_{nbn} \propto J_{cdw}$, which characterizes the CDW current oscillations, locks onto the external frequency ω_{ex} with $\omega_{nbn}/\omega_{ex} = p/q$, p and q integers. The mode-locked state is the analog of the pinned state, with the CDW remaining locked at the same frequency for a finite range of E_{dc} . This defines a mode-locked step over which the average CDW current remains fixed. Upon unlocking at lower and higher “thresholds” the current varies continuously with E_{dc} .

Over a range of temperatures the depinning and mode-locking transitions are smooth and can be analyzed in terms of dynamical critical phenomena [2]. However, at lower temperatures, CDWs also display discontinuous and sometimes hysteretic transitions. In the case of the depinning transition, hysteresis, or “switching,” is observed at a second, higher threshold field E_{th}^* [4]. More recently, Higgins *et al.* [5] observed hysteresis on mode-locked steps. Most notably, the size of the hysteresis loops on opposite edges of a given step exhibit a marked asymmetry: for certain values of the ac-drive the right edge of the step is hysteretic while the left edge displays a smooth mode-locking transition. The observed hysteresis is small in size and occurs in a parameter regime in which the sample is not switching in the absence of ac-drive. In a different set of experiments, Levy *et al.* [6] pointed to the chaotic behavior on the mode-locked steps in switching samples, in which case no well-defined mode-locking hysteresis loop can be identified. Instead, a period-doubling route to chaos with an associated low-dimensional attractor was observed while at lower frequencies high dimensional and noisy dynamics were seen.

These hysteretic and chaotic phenomena provide new challenges to our theoretical understanding of these systems. In particular, motivated by these experiments, we have studied

the mode-locking behavior of a new global coupling model [7] of CDW dynamics which has been seen previously to show some promise in explaining the temperature dependent behavior of CDWs by producing hysteresis at the depinning transition as well as delayed conduction [8]. In this paper we focus our attention on the one-dimensional version of this model, which we find produces some of the experimentally observed results and also allows us to make further predictions; however, the 1d model comes up short on some important aspects of the experiments as well.

Many aspects of CDW dynamics including the phenomena of mode-locking are well described by the Fukuyama-Lee-Rice (FLR) model [9]. An important property of this model is the so-called no passing rule [10] which requires that the current be a single-valued function of E_{dc} . Hence the FLR model cannot describe the low-temperature hysteresis exhibited at the depinning or mode-locking transitions. As pointed out by Littlewood [11] and recently shown by Levy *et al.* [7], the role of normal carriers excited across the CDW gap (Δ_{CDW}) becomes increasingly important at low temperatures. This adds a new global coupling term to the FLR equations of motion, altering the I-V characteristics significantly.

The global coupling model is obtained by incorporating current conservation within the FLR equation of motion. Ignoring the displacement current, the conservation equation is readily integrated in 1d [7] leading to the following equation suitable for numerical simulations,

$$(\gamma_0 + \gamma_1)\dot{\phi}_i = k \Delta^2 \phi_i - \sin(2\pi(\phi_i + \beta_i)) + E(t) + \frac{\gamma_1}{N} \sum_{i=1}^N \dot{\phi}_i. \quad (1)$$

Here ϕ_i represents the CDW phase at the i th impurity site, γ_0 is the time constant for CDW relaxation (we set $\gamma_0 = 1$), and γ_1 is the normal carrier resistance, assumed to have an exponential temperature-dependence: $\gamma_1 \propto \exp(\Delta_{CDW}/k_B T)$. The FLR pinning parameter is k , so that the two limits $2\pi k \ll 1$ and $2\pi k \gg 1$ correspond to the strong and weak pinning limits, respectively. The $\beta_i \in [0, 1)$ are random variables due to disorder, and Δ^2 is the lattice Laplacian. The average CDW current is given by $\bar{J}_{cdw} = \langle \dot{\phi}_i \rangle_{x,t} = E_{dc} + \bar{F}_{pin}$

where the second term is the time and space averaged pinning force

$$\bar{F}_{pin} = - \langle \sin(2\pi(\phi_i + \beta_i)) \rangle_{x,t} . \quad (2)$$

We simulate Eq. (1) with $E(t) = E_{dc} + E_{ac} \cos(\omega_{ex}t)$ using periodic boundary conditions in one dimension. Fifth-order Cash-Karp Runge-Kutta [13] is used to integrate the equations for N varying from 10 to 1024 with various values of k .

When $\gamma_1 = 0$ the standard FLR model is recovered. For non-zero γ_1 the time constant for $\dot{\phi}_i$ is adjusted and the global coupling term is relevant. It was shown [7] that the additional term gives rise to hysteresis in the regular ($E_{ac} = 0$) I-V curve; however, the size of hysteresis can be very small for small γ_1 , with the difference between the pinning and depinning fields scaling roughly as γ_1^2 for $\gamma_1 \ll 1$ [12]. For the mode-locking transition, we define the size of hysteresis S_{hyst} to be the magnitude of the difference between the locking and unlocking fields, and examine variations in S_{hyst} in different regimes.

Figure 1 shows mode-locking ($E_{ac} \neq 0$) I-V curves for different values of γ_1 in the weak pinning regime ($k = 1.0$). For $\gamma_1 = 0.0$ harmonic ($q = 1$) and many high order subharmonic ($q \neq 1$) steps are produced. As γ_1 increases hysteresis emerges asymmetrically on harmonic and some subharmonic steps. With increasing γ_1 subharmonic steps become smaller as harmonic steps grow wider. The left edges of the steps become rounded as the right edges become sharp and hysteretic. No chaotic behavior is observed. Note that within a hysteresis loop (e.g. right edge of the 0/1 step in Fig. 1(b)) two distinct mode-locked states, i.e. different p/q ratios, sometimes exist for identical set of parameters, one on the upper branch and one on the lower branch. This means that nondegenerate mode-locked attractors exist for this model, where the FLR model only exhibits degenerate mode-locked attractors displaying the same velocity [10]. We emphasize that in order to obtain this much structure in the I-V curves it was necessary to go to relatively large values of k . In comparison, while the strong pinning case of $k = 0.01$ also displays asymmetric hysteresis on mode-locked steps, the I-V structure was much simpler, displaying harmonic and only $p/2$ subharmonic steps even for large system sizes.

In analogy with the behavior for the depinning transition [7], we expect the S_{hyst} to increase with increasing γ_1 . This is always observed for strong pinning, and in general this characterizes the basic trends. However, in the weak pinning case, there are sometimes exceptions occurring in cases where we observe subharmonic steps on the upper branch of the hysteresis loops. These additional steps within a hysteresis loop appear to increase S_{hyst} as is seen by comparing Fig. 1(b) and 1(c). This feature disappears for small k or large γ_1 .

In general, the disappearance of subharmonic steps throughout the I-V curve with increasing γ_1 is consistent with experiments [6]. In the model internal deformations of the CDW must be accompanied by the screening currents of quasi-particles in order to conserve charge locally. With decreasing temperature quasi-particle conductivity ($1/\gamma_1$) decreases exponentially, inhibiting the “backflow” current. This freezes out the internal CDW deformations and results in a rigid CDW. It is well known that subharmonic mode-locking is a feature of many degrees-of-freedom dynamics and that this behavior disappears in single degree-of-freedom models [3]. Here, the disappearance is due to low dimensional behavior in a high dimensional model, arising because of the dominance of the global coupling.

Figure 2 shows the “phase diagram” in the weak pinning ($k = 1.0$) regime for $\gamma_1 = 2.5$. Figure 2 differs from the standard phase diagrams where the step width is plotted in the E_{ac} vs. E_{dc} plane. Here, in order to illustrate the existence and variations of hysteresis, we plot the full I-V curves for different values of E_{ac} . For each mode-locked state the width of the step oscillates as E_{ac} is varied. Within the FLR model, in the high velocity limit, the oscillations in harmonic mode-locked steps are described by Bessel functions [14]. In Fig. 2 the dashed lines connecting the unlocking transition points appear to be at least qualitatively consistent with the FLR case though we have not investigated the functional form in detail. The hysteresis size asymmetry is most pronounced in the first half of the first oscillations (low E_{ac}) where large hysteresis is observed. For larger E_{ac} , as in the second oscillation of the 1/1 step, hysteresis becomes much smaller and symmetric on the opposite edges of the steps. The asymmetry occurs even for high velocity mode-locked steps, e.g. 4/1 step (not shown here), at low E_{ac} values. Therefore, asymmetry appears to be a low E_{ac}

but not necessarily a low velocity phenomenon.

We suggest that an important variable governing the size of hysteresis within a given phase diagram is the pinning force at the edges of the mode-locked step. In Fig. 3 we show a schematic asymmetric mode-locked step labeling important quantities. By drawing the $\bar{J}_{cdw} = E_{dc}$ line through the step, one can visualize the magnitude of the average pinning force, $|\bar{F}_{pin}|$, as the height difference between the two curves. One might naively expect that in the middle of the step the pinning force would be zero so that at the edges of the step the pinning force would take on some extremum values as allowed by the value of E_{ac} . This is indeed the case for large E_{ac} ; the second oscillations, for example, exhibit steps for which the value of the pinning force at the center of the step is close to zero, and the value of $\bar{F}_{pin}^{edge}$ is equal in magnitude, with opposite sign, at the two edges. However, within the first oscillation, the center of the steps does not occur at $E_{dc} = \bar{J}_{cdw}^{p/q}$ i.e. where $\bar{F}_{pin} = 0$. The centers of these steps lie at higher values of E_{dc} (nonzero pinning force) and this ultimately gives rise to the hysteresis asymmetry of the steps. The center of these steps occurs at higher values of E_{dc} since for small E_{ac} the pinning force cannot change appreciably from its value when $E_{ac} = 0$. To increase $\bar{J}_{cdw}$ to the mode-locked value, $\bar{J}_{cdw}^{p/q}$, one must go to higher E_{dc} . Therefore, for low E_{ac} the steps are “shifted” to the right as is apparent from Fig. 2.

The above argument suggests that the 0/1 step should always be very nearly symmetric, since for the 0/1 step the pinning force is zero at $E_{dc} = 0$, which is always close to the center of the step [15]. Higgins *et al.* [5] did not look at the left edge of the 0/1 step so this remains to be verified. However, they did discover that for steps other than 0/1, hysteresis occurred for those values of E_{ac} where the steps were the widest, i.e. middle of the first oscillation. This is consistent with Fig. 2 where the largest hysteresis occurs in the middle of the first oscillation. More generally, the wider steps tend to have a larger pinning force at the edges, and the larger the pinning force the larger the size of hysteresis.

The observed qualitative trends in the relationship between the size of the hysteresis (e.g. $S_{hyst} = E_{T_2} - E_{T_3}$ for the right edge hysteresis in Fig. 3) and the magnitude of the pinning force (Eq. (2)) suggests that it may be worth searching for some quantitative scaling

relationship. Restricting our attention to a particular phase diagram in the strong pinning regime ($k = 0.01$ and $\gamma_1 = 1.0$, and $w_{ex} = 2\pi$), we have attempted to find such a relationship with limited success. In particular, a scaling fit between S_{hyst} and $\bar{F}_{pin}^{edge}$ appears to hold well for the symmetric steps. However, an extrapolation of this scaling breaks down in the low E_{ac} limit where the asymmetry is most pronounced. In that case on the right edges of the steps, S_{hyst} is even larger than what the extrapolation predicts, while S_{hyst} is smaller (often zero) on the left edges. This result is indicative of a crossover occurring between the high and low E_{ac} limits, which can be accounted for in a crude manner using the pinning force at the center of the step $\bar{F}_{pin}^{center}$. Note that $\bar{F}_{pin}^{center}$ is negative at low E_{ac} increasing to zero at high E_{ac} . We construct an adjusted pinning force $\bar{F}_{pin}^{adj} = \bar{F}_{pin}^{edge} + \alpha \times \bar{F}_{pin}^{center}$. Note that $\bar{F}_{pin}^{adj} = \bar{F}_{pin}^{edge}$ for symmetric steps. However, for asymmetric steps with $\alpha > 0$, $|\bar{F}_{pin}^{adj}|$ is increased on the right edge and decreased on the left edge as compared to $\bar{F}_{pin}^{edge}$, yielding a more satisfactory fit to the data. The result is shown in Fig. 4 for the 2/1 and 1/1 steps where $S_{hyst} \sim |\bar{F}_{pin}^{adj}|^\lambda$ roughly describes the data. Here, instead of fitting the data to find the best values of α and λ we have plotted the result for $\alpha = 5.0$.

Finally, regarding the discrepancies between our results and those obtained experimentally, we note that in all the simulations, for each given “sample”, the largest hysteresis typically occurs when $E_{ac} = 0$; in other words, mode-locking hysteresis occurred only in “switching samples.” Furthermore, no chaos was observed for a wide range of parameter values. This is troublesome in the light of the experiments which show hysteresis in non-switching samples, and chaos at the step edges of switching samples. One possible resolution of this may come from further study of the depinning transition. Real CDWs exhibit two transitions, only one of which becomes hysteretic at low temperatures [4]. The 1d model studied here exhibits one transition where a small hysteresis grows to a large one as a function of γ_1 . However, there is evidence [16] that a 2d generalization of the model exhibits depinning transitions which more closely resemble the experiments. In that case a new “creep” motion arises between the first and second transitions, and perhaps this qualitatively different behavior will bare out these shortcomings.

In summary, we have studied the response of the global coupling model to a joint ac+dc field. Unlike the FLR model, this model exhibits hysteresis and nondegenerate mode-locked attractors. The model produces hysteresis on mode-locked steps exhibiting size asymmetry in the low ac-field regime. The pinning force and the step offset are the primary variables which control mode-locking hysteresis within a given phase diagram. Increasing rigidity of the CDW at large values of global coupling leads to low dimensional behavior and thus vanishing of subharmonic steps. While the model exhibits no chaos in the regime studied, this low dimensional behavior may have some bearing on the low-dimensional chaotic attractors observed experimentally [6].

It is a pleasure to acknowledge useful conversations with G. Fiddich, M. Kvale, J. McCarten, C. Myers, and M.S. Sherwin. A.M. acknowledges J. Cleveland for help with graphics. Computations were performed in part at the San Diego Supercomputer Center. This work is supported by fellowships from the Alfred P. Sloan Foundation, the David and Lucile Packard Foundation, and NSF grants DMR-9212396 and DMR-9214995.

REFERENCES

- [1] For a review of CDW dynamics see *Charge-Density Waves in Solids*, No. 25 in *Modern Problems in Condensed Matter Sciences*, edited by L.P. Gorkov and G. Grüner (North-Holland, Amsterdam, 1989); G. Grüner, Rev.Mod.Phys. **60**, 1129 (1988).
- [2] D.S. Fisher, Phys.Rev.Lett. **50**, 1486 (1983); O. Narayan and D.S. Fisher, Phys.Rev.Lett. **68**, 3615 (1992).
- [3] For a review of mode-locking in CDWs see S.E. Brown and A. Zettl in ref. [1] p.223. More recent work include, J. Levy M.S. Sherwin, Phys.Rev.Lett. **67**, 2846 (1991); and A.A. Middleton, O. Biham, P. Sibani and P.B. Littlewood, Phys.Rev.Lett. **68**, 1586 (1991).
- [4] T.L. Adelman, J. McCarten, M.P. Maher, D.A. DiCarlo, and R.E. Thorne, Phys.Rev. **B 47**, 4033 (1993).
- [5] M.J. Higgins, A.A. Middleton and S. Bhattacharya, Phys.Rev.Lett. **70**, 3784 (1993).
- [6] J. Levy, M.S. Sherwin, and J. Theiler, Phys.Rev.Lett **70**, 2957 (1993).
- [7] J. Levy, M.S. Sherwin, F.F. Abraham, and K. Wiesenfeld, Phys.Rev.Lett. **68**, 2968 (1992).
- [8] J. Levy and M.S. Sherwin, submitted to Phys.Rev. **B**.
- [9] H. Fukuyama and P.A. Lee, Phys.Rev. **B 17**, 535 (1978); P.A. Lee and T.M. Rice, *ibid* **19**, 3870 (1979).
- [10] A.A. Middleton, Phys.Rev.Lett. **68**, 670 (1992).
- [11] P.B. Littlewood, Solid State Comm. **65**, 1347 (1988).
- [12] P. Colet and K.Wiesenfeld have shown that, within certain approximations, the difference in the fields $E_{th2}^* - E_{th1}^* = 1 - (1 + 2\gamma_1)^{1/2}/(1 + \gamma_1)$ (private communication).

- [13] *Numerical Recipes, The Art of Scientific Computing*, W.H. Press, B.P. Flannery, S.A. Teukolsky, and W.T. Vetterling (Cambridge University Press, Cambridge, 1992) 2nd Ed.
- [14] L. Sneddon, M.C. Cross and D.S. Fisher, Phys.Rev.Lett. **49**, 292 (1982); also see J.P. McCarten Ph.D. thesis, Cornell University (1993).
- [15] While simulations reveal that the 0/1 step is nearly symmetric, this symmetry is not exact. Note that Eq. (1) is not symmetric under $\phi_i \rightarrow -\phi_i$ and $E(t) \rightarrow -E(t)$ since the pinning term breaks the symmetry.
- [16] M. Kvale has reported 2d simulations of a model based on Ref. [11], where in addition to global coupling there is also a term associated with displacement current (unpublished).

FIGURES

FIG. 1. Mode-locking I-V curves for different values of the global coupling parameter, γ_1 . Note that (a) is the FLR limit. The curves are shifted vertically for clarity. Some steps are labeled by their p/q values in (a). The emergence of asymmetric hysteresis with increasing global coupling is evident.

FIG. 2. Mode-locking “phase diagram”; I-V curves for different values of E_{ac} . Here $k = 1.0, N = 256, \gamma_1 = 2.5$ and $\omega_{ex} = 3.7$. Curves are shifted vertically for clarity. The lowest curve is the regular I-V curve. Dashed lines connect the unlocking transition points for each step. Hysteresis varies in size for each step and is largest in the middle of the first oscillation. Note that the steps are offset to the right for low values of E_{ac} and that the offset decreases as E_{ac} increases.

FIG. 3. Geometric representation of $\bar{J}_{cdw} = E_{dc} + \bar{F}_{pin}$ on an asymmetric mode-locked step. The step is drawn schematically to label important quantities and illustrate the nature of the asymmetry.

FIG. 4. Hysteresis size vs. adjusted pinning force for 1/1 and 2/1 steps in the strong pinning regime. $S_{hyst} \sim |\bar{F}_{pin}^{edge} + \alpha \times \bar{F}_{pin}^{center}|^\lambda$ with $\alpha = 5.0$ (see text).